

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTADEG1625

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	OCT 02	CA/CAPplus enhanced with pre-1907 records from Chemisches Zentralblatt
NEWS	3	OCT 19	BEILSTEIN updated with new compounds
NEWS	4	NOV 15	Derwent Indian patent publication number format enhanced
NEWS	5	NOV 19	WPIX enhanced with XML display format
NEWS	6	NOV 30	ICSD reloaded with enhancements
NEWS	7	DEC 04	LINPADOCDB now available on STN
NEWS	8	DEC 14	BEILSTEIN pricing structure to change
NEWS	9	DEC 17	USPATOLD added to additional database clusters
NEWS	10	DEC 17	IMSDRUGCONF removed from database clusters and STN
NEWS	11	DEC 17	DGENE now includes more than 10 million sequences
NEWS	12	DEC 17	TOXCENTER enhanced with 2008 MeSH vocabulary in MEDLINE segment
NEWS	13	DEC 17	MEDLINE and LMEDLINE updated with 2008 MeSH vocabulary
NEWS	14	DEC 17	CA/CAPplus enhanced with new custom IPC display formats
NEWS	15	DEC 17	STN Viewer enhanced with full-text patent content from USPATOLD
NEWS	16	JAN 02	STN pricing information for 2008 now available
NEWS	17	JAN 16	CAS patent coverage enhanced to include exemplified prophetic substances
NEWS	18	JAN 28	USPATFULL, USPAT2, and USPATOLD enhanced with new custom IPC display formats
NEWS	19	JAN 28	MARPAT searching enhanced
NEWS	20	JAN 28	USGENE now provides USPTO sequence data within 3 days of publication
NEWS	21	JAN 28	TOXCENTER enhanced with reloaded MEDLINE segment
NEWS	22	JAN 28	MEDLINE and LMEDLINE reloaded with enhancements
NEWS	23	FEB 08	STN Express, Version 8.3, now available
NEWS	24	FEB 20	PCI now available as a replacement to DPCI
NEWS	25	FEB 25	IFIREF reloaded with enhancements
NEWS	26	FEB 25	IMSPRODUCT reloaded with enhancements
NEWS	27	FEB 29	WPINDEX/WPIDS/WPIX enhanced with ECLA and current U.S. National Patent Classification

NEWS EXPRESS FEBRUARY 08 CURRENT WINDOWS VERSION IS V8.3,
AND CURRENT DISCOVER FILE IS DATED 20 FEBRUARY 2008

NEWS HOURS	STN Operating Hours Plus Help Desk Availability
NEWS LOGIN	Welcome Banner and News Items
NEWS IPC8	For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 18:09:13 ON 19 MAR 2008

=> file caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'CAPLUS' ENTERED AT 18:09:27 ON 19 MAR 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Mar 2008 VOL 148 ISS 12

FILE LAST UPDATED: 18 Mar 2008 (20080318/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s epoxidation and catayst

15095 EPOXIDATION
249 EPOXIDATIONS
15129 EPOXIDATION
(EPOXIDATION OR EPOXIDATIONS)
26769 EPOXIDN
582 EPOXIDNS
26860 EPOXIDN
(EPOXIDN OR EPOXIDNS)
28634 EPOXIDATION
(EPOXIDATION OR EPOXIDN)
12 CATAYST
15 CATAYSTS
27 CATAYST
(CATAYST OR CATAYSTS)
L1 1 EPOXIDATION AND CATAYST

=> s epoxidation and catalyst

15095 EPOXIDATION
249 EPOXIDATIONS
15129 EPOXIDATION
(EPOXIDATION OR EPOXIDATIONS)
26769 EPOXIDN

```

        582 EPOXIDNS
        26860 EPOXIDN
              (EPOXIDN OR EPOXIDNS)
        28634 EPOXIDATION
              (EPOXIDATION OR EPOXIDN)
        794610 CATALYST
        791247 CATALYSTS
        1016631 CATALYST
              (CATALYST OR CATALYSTS)
L2      10056 EPOXIDATION AND CATALYST

```

```

=> s 12 and "Ti-MWW"
        369298 "TI"
          1300 "TIS"
        370270 "TI"
              ("TI" OR "TIS")
          217 "MWW"
            1 "MWWS"
          218 "MWW"
              ("MWW" OR "MWWS")
          40 "TI-MWW"
              ("TI" (W) "MWW")
L3      28 L2 AND "TI-MWW"

```

```

=> s 13 and precursor
        275300 PRECURSOR
        129461 PRECURSORS
        363027 PRECURSOR
              (PRECURSOR OR PRECURSORS)
L4      7 L3 AND PRECURSOR

```

```

=> s 14 and hydrothermal
        75773 HYDROTHERMAL
          5 HYDROTHERMALS
        75776 HYDROTHERMAL
              (HYDROTHERMAL OR HYDROTHERMALS)
L5      4 L4 AND HYDROTHERMAL

```

```

=> d 15 1-4 abs ibib

```

L5 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2008 ACS on STN

AB Ti-MWW was prepared by acid treatment and calcination on a lamellar precursor which was synthesized under dynamic hydrothermal crystallization conditions using piperidine as a template, and boric acid as a structure-supporting agent. The crystallization time is shortened by adding Ti-MWW seeds and increasing stirring rate. UV-visible spectra show that the precursor contains tetrahedral and octahedral Ti species. Calcination of the precursor results in a partial condensation of the octahedral Ti to form the anatase phase. The octahedral Ti species belong to a kind of extra framework Ti, which can be removed easily by the acid treatment in a short time. The calcined Ti-MWW contains the anatase when the acid-treated time is too long. The characteristic IR band at 960 cm⁻¹ is not observed for the Ti-MWW precursor, but it appears in the samples treated by acid. Epoxidn. of allyl alc. with H₂O₂ on Ti-MWW was studied. Ti-MWW with Si/Ti ratio of 20 and acid treatment for 12 .apprx. 16 h is more effective. The conversation of allyl alc. is 88.7%, and the selectivity for glycidol is 99% under the reaction conditions of 333 K and 30 min, but the catalytic activity of Ti-MWW decreases when the acid-treated time is too long.

ACCESSION NUMBER: 2006:397985 CAPLUS

DOCUMENT NUMBER: 146:69422
TITLE: Epoxidation of allyl alcohol to glycidol on
Ti-MWW molecular sieves
AUTHOR(S): Chen, Xiaohui; Fan, Zhiyong; Quan, Xia; Wei, Kemei
CORPORATE SOURCE: National Engineering Research Center of Chemical
Fertilizer Catalyst, College of Chemistry and Chemical
Engineering, Fuzhou University, Fuzhou, 350002, Peop.
Rep. China
SOURCE: Cuihua Xuebao (2006), 27(3), 285-290
CODEN: THHPD3; ISSN: 0253-9837
PUBLISHER: Kexue Chubanshe
DOCUMENT TYPE: Journal
LANGUAGE: Chinese

L5 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2008 ACS on STN

AB A novel post-synthesis method has been proposed to prepare a highly efficient titanosilicate catalyst with the MWW topol., Ti-MWW. The so-called reversible structural conversion method is based on a structural interchange between 3-dimensional MWW silicate and its lamellar precursor together a simultaneous incorporation of titanium through the treatment with an aqueous solution of titanium source and cyclic amine. This method overcomes the restriction problem between pore windows and titanium precursor probably encountered in the usual method using TiCl₄ vapor at elevated temps. The catalytic properties of postsynthesized Ti-MWW have been investigated by comparing with hydrothermally synthesized one as well as the conventional titanosilicates of TS-1 and Ti-Beta in the epoxidn. of various alkenes with hydrogen peroxide. Ti-MWW proves to be more effective in the epoxidn. of both simple and functionalized alkenes. Particularly, the novel postsynthesis method of a reversible structural conversion incorporates more active Ti species into the framework than the conventional hydrothermal synthesis, leading to the most active epoxidn. titanosilicate catalyst so far. Ti-MWW has been further delaminated into thin sheet material which possesses an extremely open and accessible surface area but maintains the basic structure of zeolite. Delaminated Ti-MWW catalyzes the epoxidn. of bulky substrates of various cycloalkenes more actively than other titanosilicates including mesoporous Ti-MCM-41.

ACCESSION NUMBER: 2005:418382 CAPLUS
DOCUMENT NUMBER: 144:150676
TITLE: MWW-type titanosilicate: novel preparation and high efficiency in the epoxidation of various alkenes
AUTHOR(S): Wu, P.; Fan, W.; Nuntasri, D.; Tatsumi, T.
CORPORATE SOURCE: Graduate School of Engineering, Yokohama National University, Hodogayaku, Yokohama, 2408501, Japan
SOURCE: Studies in Surface Science and Catalysis (2004), 154C(Recent Advances in the Science and Technology of Zeolites and Related Materials), 2581-2588
CODEN: SSCTDM; ISSN: 0167-2991
PUBLISHER: Elsevier B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2008 ACS on STN

AB MWW type titanosilicate, Ti-MWW, has been synthesized by the dry-gel conversion (DGC) method, and its physicochem. properties and catalytic performance in the liquid-phase epoxidn. of alkene have been compared with that of hydrothermally synthesized (HTS)

Ti-MWW. The roles in the crystallization of silica source, alkali cation, cyclic amine as a structure-directing agent (SDA), and boric acid structure-supporting agent have been investigated. The crystallization of Ti-MWW did not occur for the dry gels free of boric acid, but was feasible at a Si/B molar ratio as high as 12 in marked contrast to the ratio of 0.75 required in the hydrothermal synthesis. The sodium as a mineralization agent was not necessary and on the contrary inhibited the crystallization particularly at a high content. The seeding technique using deboronated MWW effectively accelerated the crystallization speed and reduced the amount of boric acid required.

As-synthesized

Ti-MWW-DGC lamellar precursors contained both tetrahedral and octahedral species but the latter was selectively removed by acid treatment. Ti-MWW-DGC catalysts showed lower intrinsic activity than Ti-MWW-HTS in the epoxidn. of hex-1-ene with hydrogen peroxide probably because the crystal size of the former was 10-20 times as large as that of the latter and then imposed significant diffusion problems for both the substrates and the products.

ACCESSION NUMBER: 2005:224837 CAPLUS
DOCUMENT NUMBER: 142:323629
TITLE: Synthesis of Ti-MWW by a dry-gel conversion method
AUTHOR(S): Wu, Peng; Miyaji, Takayuki; Liu, Yueming; He, Minyuan; Tatsumi, Takashi
CORPORATE SOURCE: Shanghai Key Laboratory of Green Chemistry and Chemical Processes, Department of Chemistry, East China Normal University, Shanghai, 200062, Peop. Rep. China
SOURCE: Catalysis Today (2005), 99(1-2), 233-240
CODEN: CATTEA; ISSN: 0920-5861
PUBLISHER: Elsevier B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2008 ACS on STN

AB A novel titanosilicate with the MWW topol., Ti-MWW, has been prepared by direct hydrothermal synthesis using boric acid as a structure-supporting agent, and also by post-incorporation of tetrahedral Ti species into MWW silicalite through controlled structural conversions between three-dimensional crystalline silicalite and the lamellar precursor. Ti-MWW is further converted by delamination into a thin sheet material applicable to the reaction of bulky reactants. Both direct hydrothermal synthesis and postsynthesis methods make it possible to introduce a controllable amount of Ti species into the MWW structure. An acid treatment of uncalcined samples is essentially important for the removal of the extraframework octahedral Ti species located on the exterior layer surface. The catalytic properties of Ti-MWW have been compared with those of conventional titanosilicates (TS-1, TS-2, Ti-Beta, Ti-MOR, Ti-MCM-41, etc.) in the epoxidn. of various alkenes with hydrogen peroxide. Hydrothermally synthesized Ti-MWW proves to be more effective in the epoxidn. of linear alkenes including functionalized ones, and also exhibits considerable activity for cycloalkenes. Moreover, it shows a unique shape selectivity not shared with other titanosilicates in the epoxidn. of cis/trans geometric alkene isomers. Postsynthesized Ti-MWW, nearly free of boron, catalyzes the alkene epoxidn. more effectively as a result of the tetrahedral Ti species different from those

resulting from the direct synthesis, which turns out to be the most active epoxidn. titanasilicate catalyst so far. Delaminated Ti-MWW, possessing an extremely open and accessible surface area but maintaining the basic structure of zeolite, catalyzes the epoxidn. of various cycloalkenes more actively than large pore titanosilicates including mesoporous Ti-MCM-41.

ACCESSION NUMBER: 2004:378219 CAPLUS
DOCUMENT NUMBER: 141:213439
TITLE: A New Generation of Titanosilicate Catalyst:
Preparation and Application to Liquid-Phase
Epoxidation of Alkenes
AUTHOR(S): Wu, Peng; Tatsumi, Takashi
CORPORATE SOURCE: Department of Chemistry, Shanghai Key Laboratory of
Green Chemistry and Chemical Processes, East China
Normal University, Shanghai, 200062, Peop. Rep. China
SOURCE: Catalysis Surveys from Asia (2004), 8(2), 137-148
CODEN: CSAABF; ISSN: 1571-1013
PUBLISHER: Kluwer Academic/Plenum Publishers
DOCUMENT TYPE: Journal
LANGUAGE: English
REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> FIL STNGUIDE
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
31.48	31.69

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-3.20	-3.20

CA SUBSCRIBER PRICE

FILE 'STNGUIDE' ENTERED AT 18:12:53 ON 19 MAR 2008
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Mar 14, 2008 (20080314/UP).

=> log off
ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF
LOGOFF? (Y)/N/HOLD:y
STN INTERNATIONAL LOGOFF AT 18:25:17 ON 19 MAR 2008